

PCTWORLD INTELLECTUAL
PROPERTY ORGANIZATION

WO 9602244A1

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/19, 31/20, 31/225, 31/23, 31/70		A1	(43) International Publication Date: 1 February 1996 (01.02.96)
(21) International Application Number: PCT/GB95/01696		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).	
(22) International Filing Date: 19 July 1995 (19.07.95)			
(30) Priority Data: 110380 20 July 1994 (20.07.94) IL			
(71) Applicant: AGIS INDUSTRIES (1983) LTD. [IL/IL]; 29 Lehi Street, 51200 Beni Brak (IL).			
(74) Agent: ABLEWHITE, Alan, James; Marks & Clerk, 57-60 Lincoln's Inn Fields, London WC2A 3LS (GB).			
(72) Inventors: SINTOV, Amnon; 47A Rothem Street, 84965 Omer (IL). UZAN, Rina; 38/23 Rahavat Zefat, 84770 Beer-Sheva (IL).		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(74) Agent: ABLEWHITE, Alan, James; Marks & Clerk, 57-60 Lincoln's Inn Fields, London WC2A 3LS (GB).			
(54) Title: A TOPICAL PREPARATION FOR THE PREVENTION AND TREATMENT OF LESIONS AND SORES ASSOCIATED WITH A HERPES VIRUS			
(57) Abstract			

The invention provides a topical pharmaceutical composition for the prevention and treatment of lesions and sores of the skin or mucosa associated with a herpes virus, comprising a carboxylic or dicarboxylic acid or a salt thereof as active ingredient therein, in combination with a pharmaceutically or cosmetically acceptable carrier.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**A TOPICAL PREPARATION FOR THE PREVENTION AND TREATMENT
OF LESIONS AND SORES ASSOCIATED WITH A HERPES VIRUS**

The present invention relates to a topical pharmaceutical composition and to methods for the treatment of a herpes virus using the same.

More particularly, the present invention relates to a topical composition for the prevention and treatment of lesions and sores associated with a herpes virus, said composition containing a carboxylic or dicarboxylic acid or a salt thereof as active ingredient therein, in a pharmaceutically or cosmetically acceptable carrier.

As compared to other areas of infectious disease, antiviral therapy has rapidly advanced during the last decades. However, despite intense efforts by pharmaceutical companies, relatively few antiviral drugs are available, in contrast to the plethora of antibiotics in clinical use. There are two main reasons for the limited use of antiviral agents: (1) most drug candidates have proven too toxic in humans, and (2) the widespread use of some antiviral agents has led to emergence of drug-resistant virus strains, particularly in immunocompromised patients.

Due to the above constraints in use of antiviral agents, and following surprising results obtained and presented herewith, the present invention is meant to provide novel compositions which prevent, and/or provide, symptomatic relief of lesions and sores associated with a herpes virus, and which compositions do not require the presence of a nucleoside analogue or any other antiviral agent now in use.

- 2 -

More specifically, in Israel Specification 104,283 there is described and claimed an antiviral topical pharmaceutical composition for treating viral diseases of the skin or mucosa, comprising a poorly soluble antiviral nucleoside derivative dispersed in an aqueous gel carrier containing a gelling agent and a water-solubilized carboxylic or dicarboxylic acid salt. The teachings of said specification are incorporated herein by reference.

The invention described in said specification was based on the discovery that said formulation achieved better anti-viral effect than the Zovirax[®] ointment or cream available on the market today.

After further research and development, however, it was surprisingly discovered that the composition described in said specification, in what was thought to be a placebo and in which the soluble anti-viral nucleoside derivative, i.e., acyclovir, was absent, also achieved better effect than the Zovirax[®] ointment or cream available on the market today.

Therefore, in light of this surprising discovery, the present invention relates to a topical pharmaceutical composition for the prevention and treatment of lesions and sores of the skin or mucosa associated with a herpes virus, comprising a carboxylic or dicarboxylic acid or a salt thereof as active ingredient therein, in combination with a pharmaceutically or cosmetically acceptable carrier.

More specifically, the present invention provides a topical pharmaceutical composition wherein said salt is selected from the group consisting of linoleates, elaidates, palmitates, myristates, oleates, malonates, succinates, adipates, pimelates, maleates, fumarates or azelates.

- 3 -

Especially preferred for use in the present invention is a water-solubilized C_{16} - C_{18} carboxylic acid salt, such as an alkali oleate.

Preferably, said topical composition will comprise a polyhydroxy compound selected from the group consisting of glycerine, propylene glycol, and polyethylene glycol.

Thus, the present invention, in especially preferred embodiments, provides a composition containing between about 0.1% and about 30% of a combination of alkali oleate and oleic acid, about 0 to 70% propylene glycol, and a pharmaceutically acceptable carrier. The pH of said composition is preferably adjusted to between 7 and 8.

The invention also provides a method for the prevention and treatment of lesions and sores of the skin or mucosa associated with a herpes virus, comprising administering a topical pharmaceutical composition comprising a carboxylic or dicarboxylic acid or a salt thereof as active ingredient therein, in an effective amount for the treatment of lesions and sores of the skin or mucosa, in combination with a pharmaceutically or cosmetically acceptable carrier.

Said composition can be effective for a wide range of virus-associated and viral-like diseases. These include herpes simplex labialis, post-herpetic neuralgia, recurrent genital herpes, cancer sore, aphthous stomatitis, vulvar vestibulities, etc.

While, as indicated, it has been discovered that the above composition is effective in itself, the carboxylic acid salt of the present invention can obviously also be combined in a pharmaceutical composition with an additional

- 4 -

poorly soluble antiviral nucleoside derivative, such as acyclovir, vidarabine, azidothymidine and ganciclovir.

The pharmaceutically or cosmetically acceptable vehicle utilizable in the compositions of the present invention can be selected from the group comprising an oil-in-water or water-in-oil emulsion, solution, cream, suspension, gel, aerosol, or powder.

Oil-in-water or water-in-oil emulsions are formulated in ways that a stable topical ointment, lotion, cream, stick or foam is achieved. The stabilization of the topical emulsions may be established and optimized by using the preferred combinations of hydrophilic and lipophilic emulsifiers, properly aligned at the water/oil interface. The emulsifying agents and their concentrations and proportions may be chosen according to the principle of the well-established HLB method published by W.C. Griffen ["H.L.B. - The Hydrophilic-Lipophilic Balance," J. Soc. Cos. Met. Chem., Vol. 5, p. 249 (1954)]].

In the case where the composition according to the invention is an emulsion, the oil phase is selected from the group consisting of beeswax, spermaceti, 2-octyl dodecanol, lanolin, sodium C₁₂₋₁₅ alcohols sulphate, esters of fatty acids and high molecular weight alcohols such as cetyl palmitate and cetearyl octanoate, esters of fatty acids and branched alcohols or polyols such as isopropyl palmitate or myristate, cocoglycerides, cosbiol, wool alcohols, cocoa butter, stearyl alcohol, cholesterol, liquid paraffin, soft paraffin, hard paraffin, or the like.

The emulsifying agents used for the purpose of dispersion of the above-mentioned fats or oils and the like

- 5 -

in the aqueous phase are advantageously selected from the group of non-ionic surfactants consisting of sorbitan sesquioleate, PEG-5 glyceryl stearate, poloxamers, cetostearyl alcohol, polysorbate 60, sorbitan monostearate, sorbitan monooleate, and glyceryl monostearate.

In the case where the composition according to the invention is a gel or solution, the composition preferably comprises an oleic acid/oleate salt, and generally a lower alkanol having from one to four carbon atoms, water, a gelifying agent (if a gel), one or more polyhydric alcohols selected from the group consisting of a lower alkylene glycol having from two to four carbon atoms, glycerine, and polyethylene glycol, having an average molecular weight from 200 to 2000, and a base, e.g., sodium hydroxide, or an acid, e.g., citric acid, for pH adjustment.

The gelifying agents are selected from the group consisting of polysaccharides such as cellulose derivatives, acrylic polymers, proteins, polyhydroxy compounds such as polyethylene glycol having an average molecular weight from 400 to 2000, and polyoxyethylene-3-cetylstearyl alcohol, known as Emulgin B3.

All semi-solid topical preparations should preferably be stable and consistent, non-leaky, non-staining, and non-greasy.

In the case where the composition according to the present invention is a powder, the composition preferably comprises an oleic acid and/or alkali oleate, and generally a diluting powder compound suitable as a lubricant. This lubricant is selected from the group consisting of talc, microcrystalline cellulose, polyvinyl pyrrolidone, metal

- 6 -

stearates, lactose or starch known to have non-irritating, non-toxic and inert properties.

In accordance with another aspect of the invention, the oleic acid and/or oleate salt could be topically applied in a slow-release manner using an adhesive sponge bandage, or, alternatively, a gauze or sponge sandwich containing a layer of the active principals of the invention situated between upper and lower absorbent layers.

The carboxylic/dicarboxylic acids and/or their salts of the present invention can also be applied onto the mucosa, for example, as a buccal gel or vaginal preparation. For this purpose, several bioadhesive polymers are selected from the group consisting of polyethylene glycols, cellulose derivatives, starch, and polyacrylic acid such as polycarbophil and Carbopol 934.

As described hereinbefore, the vehicles can be in the form of a cream, lotion, ointment, gel, stick, topical solution, gargle solution, foam, spray, liquid soap, or powder. From the point of view regarding the formulation characteristics, the pharmaceutical preparations could be processed as a water-in-oil or an oil-in-water emulsion, clear solution, gel solution, aerosol, powder mix, film-forming liquid, bioadhesive preparation, detergents-containing gel, suspension in gel, liquid, or emulsion, etc.

While the invention will now be described in connection with certain preferred embodiments in the following examples so that aspects thereof may be more fully understood and appreciated, it is not intended to limit the invention to these particular embodiments. On the contrary, it is intended to cover all alternatives, modifications and

- 7 -

equivalents as may be included within the scope of the invention as defined by the appended claims. Thus, the following examples which include preferred embodiments will serve to illustrate the practice of this invention, it being understood that the particulars shown are by way of example and for purposes of illustrative discussion of preferred embodiments of the present invention only and are presented in the cause of providing what is believed to be the most useful and readily understood description of formulation procedures as well as of the principles and conceptual aspects of the invention.

EXAMPLE 1

Water-in-Oil Ointment

<u>Ingredient</u>	<u>Percent by Weight</u>
DEHYMULS E	7.00
White beeswax	4.00
White vaseline	20.00
Heavy liquid Paraffin	10.26
Propylene glycol	25.00
Oleic acid	0.30
Sodium hydroxide	0.44
Purified water	q.s.
Citric acid 20% solution to pH 7.6-7.8 in aq. sol.	

DEHYMULS E = sorbitan sesquioleate (and) penta-erythrityl tetracocoate (and) stearyl citrate (and) beeswax (and) aluminum stearate.

Emulsification is effected by heating the oil phase (ingredients 1-4) to 88°C and blending while mixing into the aqueous phase, preheated to 75°C. The aqueous phase is prepared by dissolution of sodium hydroxide, oleic acid, propylene glycol by sequence in purified water, followed by adjusting the pH using citric acid solution.

- 8 -

EXAMPLE 2Oil-in-Water Cream

<u>Ingredient</u>	<u>Percent by Weight</u>
ARLATONE 983 S	6.25
CUTINA CBS	8.75
Cetearyl octanoate	3.75
Propylene glycol	38.00
Oleic acid	3.00
Sodium hydroxide	0.44
Purified water	q.s.
Citric acid 20% solution to pH 7.6-7.8 in aq. sol.	

ARLATONE 983 S = polyoxyethylene-5-glyceryl stearate.

CUTINA CBS = glyceryl stearate (and) cetearyl alcohol (and) cetyl palmitate (and) coco-glycerides.

The oil phase (ingredients 1-3) is heated to 80°C while mixing until a uniform liquid is achieved. The oil phase is then added into the pre-prepared and 80°C preheated aqueous phase (see Example 1), while mixing and homogenizing at 80°C.

- 9 -

EXAMPLE 3Oil-in-Water Cream

<u>Ingredient</u>	<u>Percent by Weight</u>
Mineral oil	5.00
LANNETE WAX SX	7.50
Vaseline	12.50
Propylene glycol	38.00
Oleic acid	3.00
Sodium hydroxide	0.44
Purified water	q.s.
Citric acid 20% solution to pH 7.6-7.8 in aq. sol.	

LANNETE WAX SX = cetearyl alcohol (and) sodium C₁₂₋₁₅
alcohols sulphate.

The oil phase (ingredients 1-3) is heated to 80°C while mixing until a uniform liquid is achieved. The oil phase is then added into the pre-prepared and 80°C preheated aqueous phase (see Example 1) while mixing and homogenizing at 80°C.

- 10 -

EXAMPLE 4Water-in-Oil Cream

<u>Ingredient</u>	<u>Percent by Weight</u>
<u>Phase A:</u>	
DRAGOSAN w/o	8.00
VESTAN-80	6.00
Mygliol-812	10.00
DRAGOXAT EH	4.00

Phase B:

Oleic acid	3.00
Propylene glycol	30.00
Sodium hydroxide	0.44
Purified water	q.s.
Citric acid 20% solution to pH 7.6-7.8 in aq. sol.	

DRAGOSAN w/o = a mixture of sorbitan isostearate, hydrogenated castor oil, ceresin, beeswax and mineral oil.

VESTAN-80 = mineral oil.

Mygliol-812 = caprylic/capric acids.

DRAGOXAT EH = octyldodecyl octanoate.

Phase A is heated while stirring at 80°C. Phase B is prepared as in Example 1. At 80°C, phase B is emulsified in phase A. On cooling to 50°C, the cream is continuously mixed and homogenized.

- 11 -

EXAMPLE 5Gel Preparation

<u>Ingredient</u>	<u>Percent by Weight</u>
Oleic acid	4.40
Sodium hydroxide	0.64
Propylene glycol	50.00
Methocel K-15	1.90
Purified water	q.s.
Citric acid 20% solution to pH 7.3-7.5 in gel	

Methocel K-15 = hydroxypropyl methylcellulose

Manufacturing procedure as in preparation of aqueous phase in Example 1, followed by dissolution of the gelling agent - Methocel.

EXAMPLE 6Foam Preparation

<u>Ingredient</u>	<u>Percent by Weight</u>
CETIOL HE	9.00
Propylene glycol	38.00
Oleic acid	3.00
Sodium hydroxide	0.44
BRIJ 35	1.00
ARLACEL 186	0.10
Purified water	q.s.
Citric acid 20% solution to pH 7.6-7.8 in concentrate	
Isopropane/butane/isobutane 90:9:1	+5-7%

CETIOL HE = polyethylene glycol 7 glyceryl cocoate.

BRIJ 35 = polyoxyethylene 23 lauryl alcohol.

ARLACEL 186 = glyceryl mono and dioleate.

At ambient temperature, BRIJ 35, ARLACEL 186 and CETIOL HE are dissolved in the aqueous solution of sodium oleate

- 12 -

preprepared according to the procedure described in Example 1 (aqueous phase). The pH is adjusted as the last step, using citric acid solution. The ready concentrate is filled into aerosol containers and sealed with valves. The propellant gas is then filled under pressure.

EXAMPLE 7

Orabase (Adhesive Oral Gel)

<u>Ingredient</u>	<u>Percent by Weight</u>
Oleic acid	3.00
Sodium hydroxide	0.44
Ethyl alcohol	10.00
Propylene glycol	30.00
GANTREZ AN 169 (GAF)	2.50
Preservatives	0.12
Purified water	q.s.
Citric acid 20% solution to pH 7.6-7.8 in sol.	

GANTREZ AN 169 (GAF) = gelling agent.

Manufacturing procedure as in preparation of aqueous phase in Example 1, followed by dissolution of the gelling agent, GANTREZ.

- 13 -

EXAMPLE 8Gargle Solution

<u>Ingredient</u>	<u>Percent by Weight</u>
Oleic acid	3.00
Sodium hydroxide	0.44
Propylene glycol	38.00
Sodium saccharine	0.10
Flavour	0.10
Purified water	q.s.
Citric acid 20% solution to pH 7.6-7.8 in sol.	

Manufacturing procedure as in preparation of aqueous phase in Example 1.

EXAMPLE 9Comparative Effectiveness of Preparations

In-Vivo Preventive Testing of Gel (Example 5) in the Topical Treatment of Experimental Cutaneous Herpes Simplex Virus Type 1 (HSV-1) Infection

In a double-blind study, the efficacy of a gel (Example 5) according to the present invention was evaluated as compared to an untreated control and a drug-containing (5% acyclovir) cream, currently marketed by Burroughs-Wellcome under the trade name Zovirax[®].

A guinea pig model was selected for the cutaneous HSV-1, because the model mimics the human herpes simplex labialis infection [Hubler, et al., J. Invest. Dermatol., Vol. 62, pp. 92-95 (1974)]. Forty-eight animals (young male D. Hartly guinea pig CRL:(HA)BR strain, Charles River, U.K.) were assigned to the experiment.

- 14 -

The stock virus (HSV-1, HF strain) was grown on VERO cells at multiplicity of infection of 0.1. The virus was grown for three days at 37°C and thereafter frozen and thawed three times. Then, it was cleared by two-speed centrifugation at 15,000 rpm for 15 minutes. The stock virus was then dispensed into 1 ml vials and stored at -70°C.

The virus was inoculated intradermally on three locations on the back of each animal. The area into which the virus was to be inoculated was divided into three squares with a marking pen. In the middle of each area, about 20 microliters of HSV at a concentration of 10^6 PFU/ml was applied by 10 injections close to each other. In order to assign the location of the various treatments on each animal and to overcome any gradient effect, a latin-square design was used.

Treatment was started 2-3 hours after inoculation and consists of three daily applications of the topical preparations (about 50 mg) for a period of three days (9 treatments). Animals were inspected once daily, and the lesions or inoculated sites were scored in the following manner:

<u>Score</u>	<u>Description</u>
0.0	No signs detected
0.5	Erythema and slight edema at the site
1.0	Erythema and one or two small vesicles
2.0	Erythema and numerous small vesicles
2.5	Numerous medium-sized vesicles
3.0	Numerous large vesicles

In addition to the lesion scoring described above, the number of vesicles in each marked area was counted and recorded at the same time as the scoring.

- 15 -

Table 1 shows the mean and the standard deviation of the scores observed at days 2 and 3 from inoculation. Table 2 presents the mean vesicles quantities at days 2 and 3, and Table 3 shows the combination of the two parameters, providing a more comprehensible picture of the diseased state. The results clearly demonstrate the significant superiority of the composition of the invention over the acyclovir-containing commercial cream and the untreated control. The lowest average response at day 3 was associated with the non-drug-containing gel, followed by Zovirax[®] cream, and finally untreated control. The score and number of vesicles for the untreated control areas were almost as high as the respective values of Zovirax[®] cream, while those of the gel formulation were significantly lower. Thus, the responses (score and vesicles) to the gel were relatively low at day 3. In order to weight the evidence that this treatment was really better than that of the Zovirax[®] cream, the difference between responses to this treatment and those observed with respect to Zovirax[®] were tested for significance. The results were highly significant ($p < 0.001$) for each of the responses.

TABLE 1

Mean Score Number at Each Infection Site
(Mean \pm S.E.)

	<u>Day 2</u>	<u>Day 3</u>
Untreated	2.06 (\pm 0.11)	1.90 (\pm 0.13)
No-drug gel	1.97 (\pm 0.12)	0.97 (\pm 0.13)
Zovirax [®]	2.05 (\pm 0.09)	1.83 (\pm 0.12)

- 16 -

TABLE 2

Mean Count of Vesicles at Each Infection Site
(Mean \pm S.E.)

	<u>Day 2</u>	<u>Day 3</u>
Untreated	6.17 (\pm 0.46)	5.83 (\pm 0.48)
No-drug gel	4.40 (\pm 0.43)	1.13 (\pm 0.32)
Zovirax [®]	5.98 (\pm 0.43)	5.17 (\pm 0.46)

TABLE 3

Mean of Combined Parameters (Score X Vesicles)
at Each Infection Site
(Mean \pm S.E.)

	<u>Day 2</u>	<u>Day 3</u>
Untreated	14.48 (\pm 1.19)	13.58 (\pm 1.18)
No-drug gel	10.08 (\pm 1.02)	2.47 (\pm 0.75)
Zovirax [®]	13.40 (\pm 1.04)	11.30 (\pm 1.16)

It will be evident to those skilled in the art that the invention is not limited to the details of the foregoing illustrative examples and that the present invention may be embodied in other specific forms without departing from the essential attributes thereof, and it is therefore desired that the present embodiments and examples be considered in all respects as illustrative and not restrictive, reference being made to the appended claims, rather than to the foregoing description, and all changes which come within the meaning and range of equivalency of the claims are therefore intended to be embraced therein.

- 17 -

WHAT IS CLAIMED IS:

1. A topical pharmaceutical composition for the prevention and treatment of lesions and sores of the skin or mucosa associated with a herpes virus, comprising a carboxylic or dicarboxylic acid or a salt thereof as active ingredient therein, in combination with a pharmaceutically or cosmetically acceptable carrier.
2. A topical pharmaceutical composition according to claim 1, wherein said salt is selected from the group consisting of linoleates, elaidates, palmitates, myristates, oleates, malonates, succinates, adipates, pimelates, maleates, fumarates or azelates.
3. A topical pharmaceutical composition according to claim 1, wherein said salt is a water-solubilized C_{16} - C_{18} carboxylic acid salt.
4. A topical pharmaceutical composition according to claim 1, wherein said carboxylic acid salt is an oleate.
5. A topical pharmaceutical composition according to claim 1, further comprising a carboxylic or dicarboxylic acid in combination with said acid salt.
6. A topical pharmaceutical composition according to claim 1, further comprising a polyhydroxy compound.

- 18 -

7. A topical pharmaceutical composition according to claim 6, wherein said polyhydroxy compound is selected from the group consisting of glycerin, polypropylene glycol, and polyethylene glycol.

8. A topical pharmaceutical composition according to claim 1, wherein said carboxylic acid salt is an alkali oleate.

9. A topical pharmaceutical composition according to claim 1, further comprising an additional poorly soluble antiviral nucleoside derivative selected from the group consisting of acyclovir, vidarabine, azidothymidine and ganciclovir.

10. A topical pharmaceutical composition for the prevention and treatment of lesions and sores of the skin or mucosa associated with a herpes virus, consisting essentially of a carboxylic or dicarboxylic acid or a salt thereof as active ingredient therein, in combination with a pharmaceutically or cosmetically acceptable carrier.

- 19 -

11. A method for the prevention and treatment of lesions and sores of the skin or mucosa associated with a herpes virus, comprising administering a topical pharmaceutical composition comprising a carboxylic or dicarboxylic acid or a salt thereof as active ingredient therein, in an effective amount for the treatment of lesions and sores of the skin or mucosa, in combination with a pharmaceutically or cosmetically acceptable carrier.

12. The use of a carboxylic or dicarboxylic acid as an active curative or prophylactic agent in the preparation of a topical pharmaceutical composition for the prevention and treatment of lesions and sores of the skin or mucosa associated with a herpes virus.

13. The use according to claim 12 in the preparation of a composition according to any of claims 1 to 10.

INTERNATIONAL SEARCH REPORT

Int. l. Application No
PCT/GB 95/01696

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/19 A61K31/20 A61K31/225 A61K31/23 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO,A,94 15614 (AGIS INDUSTRIES LIMITED) 21 July 1994 see claims 1-11	1-13
X	EP,A,0 465 423 (RESEARCH FOUNDATION FOR MENTAL HYGIENE INC) 8 January 1992 * claims & p.23, 1.30; Table 2 * see the whole document	1-5, 8, 10-13
Y	---	6,7,9
X	GB,A,2 004 741 (GENIEYZ FERNAND) 11 April 1979 * p.3, 1.36; p.1, 1.24; claims * see the whole document	1-4, 10-13
Y	---	5-9
X	US,A,4 520 132 (KINSOLVING C.R.) 28 May 1985 see claims 1-13 see the whole document	1,5-7, 10-13
Y	---	2-4,8,9
	--- -/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

10 November 1995

Date of mailing of the international search report

24.11.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Uiber. P

INTERNATIONAL SEARCH REPORT

Inte onal Application No
PCT/GB 95/01696

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,4 513 008 (REVICI ET AL) 23 April 1985 * claims; col.3, 1.40-50; col.7, 1.28-29 *	1,5-7, 10-13
Y	see the whole document ---	2-4,8,9
X	WO,A,93 25213 (PITMY INTERNATIONAL N.V.) 23 December 1993 * claims 1,6; p.3, 1.8-21 *	1-7, 10-13
Y	see the whole document ---	8,9
X	WO,A,88 00465 (SCHERING AG) 28 January 1988 * Ex. 1; claims *	1,2,5-7, 10
X	CHEM. PHARM. BULL., vol. 38, no. 11, 1990 pages 3086-9, SEKI ET AL 'enhanced transdermal delivery of zidovudine in rats and human skins' * p.3807, fig.1; Abstract *	9
Y	see the whole document ---	1-8, 10-13
Y	INFECTION, vol. 18, no. 6, 1990 pages 372-5, GALDIERO ET AL 'effect of modification of HEp2 cell membrane lipidic phase on susceptibility to infection from herpes simplex virus' see the whole document ---	1-13
Y	ARCHIVES OF VIROLOGY, vol. 66, 1980 pages 301-7, KOHN ET AL 'unsaturated free fatty acids inactivate animal enveloped viruses' see the whole document ---	1-13
Y	US,A,4 910 224 (HABIB ET AL) 20 March 1990 * claims; Tables 4 and 5 *	1-13

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 95/01696

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9415614	21-07-94	AU-B- 5711894 CA-A- 2152907 EP-A- 0677126	15-08-94 21-07-94 18-10-95
EP-A-465423	08-01-92	AU-B- 7924891 CA-A- 2045262 GR-B- 1000958 JP-A- 6040810 US-A- 5434182	02-01-92 26-12-91 16-03-93 15-02-94 18-07-95
GB-A-2004741	11-04-79	LU-A- 78168 BE-A- 870647 DE-A- 2841346 FR-A, B 2403800	25-05-79 21-03-79 05-04-79 20-04-79
US-A-4520132	28-05-85	AU-B- 567508 AU-B- 1919583 CA-A- 1218608 EP-A, B 0105448 JP-A- 59070615 US-E- RE32990	26-11-87 05-04-84 03-03-87 18-04-84 21-04-84 18-07-89
US-A-4513008	23-04-85	NONE	
WO-A-9325213	23-12-93	CA-A- 2136614	23-12-93
WO-A-8800465	28-01-88	DE-A- 3623862 AU-B- 7701187 EP-A, B 0305407	21-01-88 10-02-88 08-03-89
US-A-4910224	20-03-90	AU-B- 604226 AU-B- 6872487 AU-B- 7028887 EP-A- 0238198 EP-A- 0257081 WO-A- 8704926 JP-A- 62265222 JP-T- 63502660	13-12-90 20-08-87 09-09-87 23-09-87 02-03-88 27-08-87 18-11-87 06-10-88